

Annex II: Reference publications concerning AS+MQ studies

Price RN, Nosten F, Luxemburger C, van Vugt M, Phaipun L, et al. (1997) . Artesunate/mefloquine treatment of multi-drug resistant falciparum malaria. **Thai-Burmese border**. *Trans R Soc Trop Med Hyg* 91: 574–577.

Denis MB, Tsuyuoka R, Poravuth Y, Narann TS, Seila S, Lim C, Incardona S, Lim P, Sem R, Socheat D, Christophel EM, Ringwald P. Surveillance of the efficacy of artesunate and mefloquine combination for the treatment of uncomplicated falciparum malaria in **Cambodia**. *Tropical Medicine and International Health*. 2006 Sep; 11(9): 1360-6.

Rogers WO, Sem R, Tero T, Chim P, Lim P, Muth S, Socheat D, Arie F and Wongsrichanalai C. Failure of artesunate-mefloquine combination therapy for uncomplicated *Plasmodium falciparum* malaria in southern **Cambodia**- *Malaria Journal* 2009, 8:10

Na-Bangchang K, Ruengweerayut R, Mahamad P, Ruengweerayut K, Chaijaroenkul W. Declining in efficacy of a three-day combination regimen of mefloquine-artesunate in a multi-drug resistance area along the **Thai-Myanmar** border. *Malar J*. 2010 Oct 8;9:273.

van Vugt M, Brockman A, Gemperli B, Luxemburger C, Gathmann I, et al. (1998) Randomized comparison of artemether-benflumetol and artesunate mefloquine in treatment of multidrug-resistant falciparum malaria. **Thailand**. *Antimicrob Agents Chemother* 42: 135–139.

van Vugt M, Looareesuwan S, Wilairatana P, McGready R, Villegas L, et al. (2000) Artemether-lumefantrine for the treatment of multidrug-resistant falciparum malaria. **Thailand**. *Trans R Soc Trop Med Hyg* 94: 545–548.

Lefèvre G, Looareesuwan S, Treprasertsuk S, Krudsood S, Silachamroon U, Gathmann I, Mull R, Bakshi R: A clinical and pharmacokinetic trial of six doses of artemether/lumefantrine for multidrug-resistant *Plasmodium falciparum* malaria in **Thailand**. *Am J Trop Med Hyg* 2001, 64:247-256.

van Vugt M, Leonardi E, Phaipun L, Slight T, Thway KL, et al. (2002) Treatment of uncomplicated multidrug-resistant falciparum malaria with artesunate-atovaquone-proguanil. **Thailand**. *Clin Infect Dis* 35: 1498–1504.

Suputtamongkol Y, Chindarat S, Silpasakorn S, Chaikachonpat S, Lim K, Chanthapakajee K, Kaewkaukul N, Thamlikitkul V. The efficacy of combined mefloquine-artesunate versus mefloquine-primaquine on subsequent development of *Plasmodium falciparum* gametocytemia. **Thailand**. *Am J Trop Med Hyg*. 2003 May;68(5):620-3.

Ashley EA, Krudsood S, Phaipun L, Srivilairit S, McGready R, et al. (2004) Randomized, controlled dose-optimization studies of dihydroartemisinin-piperaquine for the treatment of uncomplicated multidrug-resistant falciparum malaria in **Thailand**. *J Infect Dis* 190: 1773–1782.

Hutagalung R, Paiphun L, Ashley EA, McGready R, Brockman A, et al. (2005) A randomized trial of artemether-lumefantrine versus mefloquine-artesunate for the treatment of uncomplicated multi-drug resistant *Plasmodium falciparum* on the western border of **Thailand**. *Malar J* 4: 46.

Ashley EA, McGready R, Hutagalung R, Phaiphun L, Slight T, et al. (2005) A Randomized, Controlled Study of a Simple, Once-Daily Regimen of Dihydroartemisinin-Piperaquine for the Treatment of Uncomplicated, Multidrug-Resistant Falciparum Malaria. **Thailand**. *Clin Infect Dis* 41: 425–432.

Ashley EA, Lwin KM, McGready R, Simon WH, Phaiphun L, et al. (2006) A randomised comparative study of a new fixed combination of mefloquine artesunate compared to the separate drugs for the treatment of uncomplicated multidrug resistant falciparum malaria in **Thailand**. *Trop Med Int Health* 11: 1653–1660

Dondorp AM, Nosten F, Yi P, Das D, Phyo AP, Tarning J, Lwin KM, Arie F, Hanpithakpong W, Lee SJ, Ringwald PI, Silamut KI, Imwong M, Chotivanich K, Lim P, Herdman T, An SS, Yeung S, Singhasivanon P, Day NPJ, Lindegardh N, Socheat D and White N. Artemisinin Resistance in *Plasmodium falciparum* Malaria in **Cambodia**. *N Engl J Med* 2009;361:455-67.

Valecha N, Phyo AP, Mayxay M, Newton PN, Krudsood S, et al. (2010) An Open-Label, Randomised Study of Dihydroartemisinin-Piperaquine Versus Artesunate-Mefloquine for Falciparum Malaria in **Asia**. *PLoS ONE* 5(7): e11880.

Krudsood S Looareesuwan S, Tangpukdee N, Wilairatana P, Phumratanaprapin W, Leowattana W, Chalermrut K, Ramanathan S, Navaratnam V, Olliaro P, Vaillant M, Kiechel JR, Taylor WRJ. New fixed dose artesunate/mefloquine for treating multidrug resistant *Plasmodium falciparum* in adults – a comparative phase IIb safety and pharmacokinetic study with standard dose non-fixed artesunate plus mefloquine, **Thailand**. *Antimicrobial Agents and Chemotherapy*, Sept. 2010, p. 3730–3737

McGready R, Brockman A, Cho Thein, Cho D, van Vugt M, Luxemburger C, Chongsuphajaisiddhi, White N and Nosten F. Randomized comparison of mefloquine-artesunate versus quinine in the treatment of multidrug-resistant falciparum malaria in pregnancy. **Thailand**. *Transactions of the Royal Society of tropical Medicine and Hygiene* (2000)94,689-69

Smithuis F, van der Broek I, Katterman N, Kyaw MK, Brockman A, Lwin S, White NJ. Optimising operational use of artesunate-mefloquine: a randomised comparison of four treatment regimens. **Myanmar**. *Trans R Soc Trop Med Hyg*. 2004 Mar;98(3):182-92.

Smithuis F, Kyaw Kyaw M, Phe O, Zarli Aye K, Htet L, Barends M, Lindegardh N, Singtoroj T, Ashley E, Lwin S, Stepniewska K, White NJ. Efficacy and effectiveness of dihydroartemisinin-piperaquine versus artesunate-mefloquine in falciparum malaria: an open-label randomised comparison- **Myanmar** - *Lancet* 2006; 367: 2075–85.

Smithuis F, Moe Kyaw Kyaw, Ohn Phe, Thein Win, Pyay Phyo Aung, Aung Pyay Phyo Oo, Arkar Linn Naing, Mya Yee Nyo, Naing Zaw Htun Myint, Mallika Imwong, Ashley E, Lee SJ, White N. Effectiveness of five artemisinin combination regimens

with or without primaquine in uncomplicated falciparum malaria: an open-label randomised trial. **Myanmar**. *Lancet Infect Dis*. 2010 Oct;10(10):673-81.

Mayxay M, Khanthavong M, Lindegårdh N, Keola S, Barends M, Pongvongsa T, Yapom R, Annerberg A, Phompida S, Phetsouvanh R, White NJ, Newton PN: Randomized comparison of chloroquine plus sulfadoxine-pyrimethamine versus artesunate plus mefloquine versus artemether/lumefantrine in the treatment of uncomplicated falciparum malaria in the **Lao** People's Democratic Republic. *Clin Infect Dis* 2004, 39:1139-1147.

Stohrer JM, Dittrich S, Thongpaseuth V, Vanisaveth V, Phetsouvanh R, Phompida S, Monti F, Christophel EM, Lindegårdh N, Annerberg A, Jelinek T: Therapeutic efficacy of artemether-lumefantrine and artesunate-mefloquine for treatment of uncomplicated *Plasmodium falciparum* malaria in Luang Namtha Province, **Lao** People's Democratic Republic. *Trop Med Int Health*. 2004 Nov;9(11):1175-83.

Mayxay M, Thongpraseuth V, Khanthavong M, Lindegårdh N, Barends M, Keola S, Pongvongsa T, Phompida S, Phetsouvanh R, Stepniewska K, White N. J. and Newton P. N. An open, randomized comparison of artesunate plus mefloquine vs. dihydroartemisinin-piperaquine for the treatment of uncomplicated *Plasmodium falciparum* malaria in the **Lao** People's Democratic Republic (Laos). *Tropical Medicine and International Health*- volume 11 no 8 pp 1-9 august 2006.

Mayxay M, Keomany S, Khanthavong M, Souvannasing P, Stepniewska K, Khomthilath T, Keola S, Pongvongsa T, Phompida S, Ubben D, Valecha Neena, White NJ and Newton PN. A phase III, randomized, non-inferiority trial to assess efficacy and safety of dihydroartemisinin-piperaquine in comparison with artesunate-mefloquine in patients with uncomplicated *Plasmodium falciparum* malaria in Southern **Laos**. *Am. J. Trop. Med. Hyg.* 83 (6), 2010, pp.1221-1229

Denis MB, Tsuyuoka R, Lim P, Lindegårdh N, Yi P, Top SN, Socheat D, Fandeur T, Annerberg A, Christophel EM, Ringwald P. Efficacy of artemether-lumefantrine for the treatment of uncomplicated falciparum malaria in northwest **Cambodia**. *Tropical Medicine and International Health*. 2006 Dec; 11(12): 1800-7.

Janssens B, van Herp M, Goubert L, Chan S, Uong S, Nong S, Socheat D, Brockman A, Ashley EA and Van Damme W. A randomized open study to assess the efficacy and tolerability of dihydroartemisinin-piperaquine for the treatment of uncomplicated falciparum malaria in **Cambodia**. *Tropical Medicine and International Health*- volume 12 no 2 pp 251-259 - 2007.

Hien T T, Dolecek C, Phuong Mai P, Thi Dung N, Thanh Truong N, Hong Thai L, Thi Hoai An D, Tan Thanh T, Stepniewska K, White NJ, Farrar J. Dihydroartemisinin-piperaquine against multidrug-resistant *Plasmodium falciparum* malaria in **Vietnam**: randomised clinical trial. *Lancet* 2004; 363: 18-22.

van den Broek IV, Maung UA, Peters A, Liem L, Kamal M, Rahman M, Rahman MR, Bangali AM, Das S, Barends M, Faiz AM: Efficacy of chloroquine + sulfadoxine-pyrimethamine, mefloquine + artesunate and artemether + lumefantrine combination therapies to treat *Plasmodium falciparum* malaria in the Chittagong Hill Tracts, **Bangladesh**. *Trans R Soc Trop Med Hyg* 2005, 99:727-735.

Luxemburger C, Nosten F, Shotar, Raimond D, Chongsuphajaisiddhi T and White NJ. Oral Artesunate in the treatment of uncomplicated hyperparasitemic falciparum malaria in **Thailand**. *Am. J. Trop. Med. Hyg* 53(5) 1995, 522-525

Krudsod S, Looareesuwan S, Silachamroon U, Chalermrut K, Pittrow D, Cambon N, Mueller EA. Artesunate and mefloquine given simultaneously for three days via a prepacked blister is equally effective and tolerated as a standard sequential treatment of uncomplicated acute *Plasmodium falciparum* malaria: randomized, double-blind study in **Thailand**. *Am J Trop Med Hyg*. 2002 Nov;67(5):465-72.

Silachamroon U, Krudsod S, Thanachartwet W, Tangpukdee N, Leowattana W, Chalermrut K, Srivilairit S, Wilairatana P, Thimasarn K, Looareesuwan S. An open, randomized trial of three-day treatment with artesunate combined with a standard dose of mefloquine divided over either two or three days, for acute, uncomplicated falciparum malaria. **Thailand**. *Southeast Asian J Trop Med Public Health*. 2005 May;36(3):591-6.

Vijaykadga S, Rojanawatsirivej C, Cholpol S, Phoungmanee D, Nakavej A and Wongsrichanalai C. In vivo sensitivity monitoring of mefloquine monotherapy and artesunate-mefloquine combinations for the treatment of uncomplicated falciparum malaria in **Thailand** in 2003. *Tropical Medicine and International Health* volume 11 no 2 pp 211-219 february 2006

Krudsod S, Tangpukdee N, Thanchartwet V, Wilairatana P, Srivilairit S, Pothipak N, Jianping S, Guoqiao L, Brittenham GM, Looareesuwan S. Dose ranging studies of new artemisinin-piperaquine fixed combinations compared to standard regimens of artemisinin combination therapies for acute uncomplicated falciparum malaria. **Thailand**. *Southeast Asian J Trop Med Public Health*. 2007 Nov;38(6):971-8.

Tangpukdee N, Krudsod S, Thanachartwet V, Pengruksa C, Phophak N, Kano S, Li G, Brittenham GM, Looareesuwan S, Wilairatana P. Efficacy of Artequick versus artesunate-mefloquine in the treatment of acute uncomplicated falciparum malaria in **Thailand**. *Southeast Asian J Trop Med Public Health*. 2008 Jan;39(1):1-8.

Congpuong K, Bualombai P, Banmairuroi V, Na-Bangchang K. Compliance with a three-day course of artesunate-mefloquine combination and baseline anti-malarial treatment in an area of **Thailand** with highly multidrug resistant falciparum malaria. *Thailand. Malar J*. 2010 Feb 4;9:43.

Karbwang J, Na Bangchang K, Thanavibul A, Back DJ, Bunnag D, Harinasuta T. Pharmacokinetics of mefloquine alone or in combination with artesunate. **Thailand**. *Bull World Health Organ*. 1994;72(1):83-7.

Price RN, Luxemburger C, van Vugt M, Nosten F, Kham A, Simpson J, Looareesuwan S, Chongsuphajaisiddhi T and White NJ. Artesunate and mefloquine in the treatment of uncomplicated multidrug-resistant hyperparasitaemic falciparum malaria. **Thailand**. *Trans R Soc Trop Med Hyg*. 1998 92, 207-211

Sabchareon A, Attanath P, Chanthavanich P, Phanuaksook P, Prarinyanupharb V, Poonpanich Y, Mookmanee D, Teja-Isavadharm P, Heppner DG, Brewer TG and Chongsuphajaisiddhi. Comparative clinical trial of artesunate suppositories and oral artesunate in combination with mefloquine in the treatment of children with acute falciparum malaria. **Thailand**. *Am. J. Trop. Med. Hyg.*, 58(1), 1998, pp. 11–16

Rojanawatsirivej C, Vijaykadga S, Amklad I, Wilairatna P, Looareesuwan S. Monitoring the therapeutic efficacy of antimalarials against uncomplicated falciparum malaria in **Thailand**. *Southeast Asian J Trop Med Public Health* 2003, 34:536-541.

Hung Le Q, de Vries PJ, Binh TQ, Giao PT, Nam NV, Holman R, Kager PA. Artesunate with mefloquine at various intervals for non-severe Plasmodium falciparum malaria. **Vietnam**. *Am J Trop Med Hyg.* 2004 Aug;71(2):160-6.

Smithuis F, Shahmanesh M, Kyaw MK, Savran O, Lwin S, White NJ. Comparison of chloroquine, sulfadoxine/pyrimethamine, mefloquine and mefloquine-artesunate for the treatment of falciparum malaria in Kachin State, North **Myanmar**. *Trop Med Int Health.* 2004 Nov;9(11):1184-90.

Campbell P, Baruah S, Narain K, Rogers CC. A randomized trial comparing the efficacy of four treatment regimens for uncomplicated falciparum malaria in Assam state, **India**. *Trans R Soc Trop Med Hyg.* 2006 Feb;100(2):108-18.

Ambler MT, Dubowitz LM, Arunjerda R, Hla EP, Thwai KL, Viladpainguen J, Singhasivanon P, Luxemburger C, Nosten F, McGready R. The neurological assessment in young children treated with artesunate monotherapy or artesunate-mefloquine combination therapy for uncomplicated Plasmodium falciparum malaria. **Thailand**. *Malar J.* 2009 Sep 2;8:207.

Adam I, Aelbasit IE, Elbashir MI. Efficacies of mefloquine alone and of artesunate followed by mefloquine, for the treatment of uncomplicated, *Plasmodium falciparum* malaria in eastern **Sudan**. *Annals of Tropical Medicine and Parasitology*, Volume 99, Number 2, March 2005, pp. 111-117(7)

Faye B, Ndiaye JL, Ndiaye D, Dieng Y, Faye O and Gaye O. Efficacy and tolerability of four antimalarial combinations in the treatment of uncomplicated P falciparum malaria in **Senegal**. *Malaria Journal* 2007, 6: 80

Sagara I, Diallo A, Kone M, Coulibaly M, Diawara SI, Guindo O, Maiga H, Niamele MB, Sissoko M, Dicko A, Djimde A, Doumbo OK. A randomized trial of artesunate-mefloquine versus artemether-lumefantrine for treatment of uncomplicated Plasmodium falciparum malaria in **Mali**. *Am J Trop Med Hyg.* 2008 Nov;79(5):655-61.

Sowunmi A, Gbotosho GO, Happi C, Okuboyejo T, Folarin O, Balogun S, Michael O. Therapeutic efficacy and effects of artesunate-mefloquine and mefloquine alone on malaria-associated anemia in children with uncomplicated Plasmodium falciparum malaria in southwest **Nigeria**. *Am J Trop Med Hyg.* 2009 Dec;81(6):979-86.

Faye B, Ndiaye JL, Tine R, Sylla K, Gueye A, Lô AC, Gaye O. A randomized trial of artesunate mefloquine versus artemether lumefantrine for the treatment of uncomplicated Plasmodium falciparum malaria in **Senegalese** children. *Am J Trop Med Hyg.* 2010 Jan;82(1):140-4.

Bouyou-Akotet MK, Ramharter M, Ngoungou EB, Mamfoumbi MM, Mihindou MP, Missinou MA, Kurth F, B elard S, Agnandji ST, Issifou S, Heidecker JL, Trapp S, Kremsner PG and Kombila M. Efficacy and safety of a new pediatric artesunate mefloquine drug formulation for the treatment of uncomplicated falciparum malaria in **Gabon**. *Wien Klin Wochenschr* (2010) 122: 173–178

Whegang SY, Tahar R, Foumane VN, Soula G, Gw et H, Thalabard JC, Basco LK. Efficacy of non-artemisinin- and artemisinin-based combination therapies for uncomplicated falciparum malaria in **Cameroon**. *Malar J.* 2010 Feb 19;9:56.

Tietche F, Chelo D, Kinkela N, Ntoto M, Minjiwa Djoukoue F, Hatz C, Frey S, Frentzel A, Trapp S, Zielonka R and Mueller E. Tolerability and efficacy of a Pediatric Granule Formulation of Artesunate-Mefloquine in young children from **Cameroon** with uncomplicated falciparum malaria. *Am. J. Trop. Med. Hyg.* 82(6), 2010, pp1034-1040.

Toure OA, Kouame MG, Didier YJ, Berenger AA, Djerea K, Genevieve GO, Penali LK. Artesunate/mefloquine paediatric formulation vs. artemether/lumefantrine for the treatment of uncomplicated Plasmodium falciparum in Anonkoua kout e, **C ote d'Ivoire**. *Trop Med Int Health.* 2011 Mar;16(3):290-7.

Massougbodji A, Kone M, Kinde-Gazard D, Same-Ekobo A, Cambon N, Mueller EA. A randomized, double-blind study on the efficacy and safety of a practical three-day regimen with artesunate and mefloquine for the treatment of uncomplicated Plasmodium falciparum malaria in **Africa**. *Trans R Soc Trop Med Hyg.* 2002 Nov-Dec;96(6):655-9.

Awad MI, Alkadru AMY, Behrens RH, Baraka OZ and Eltayeb IB. Descriptive study on the efficacy and safety of Artesunate suppository in combination with other antimalarials in the treatment of severe malaria in **Sudan**. *Am. J. Trop. Med. Hyg.*, 68(2), 2003, pp. 153–158

Bhatt KM, Samia BM, Bhatt SM, Wasunna KM. Efficacy and safety of an artesunate/mefloquine combination, artequin, in the treatment of uncomplicated P. falciparum malaria in **Kenya**. *East Afr Med J.* 2006 May;83(5):236-42.

Ramharter M, Kurth FM, B elard S, Bouyou-Akotet MK, Mamfoumbi MM, Agnandji ST, Missinou MA, Adegnika AA, Issifou S, Cambon N, Heidecker JL, Kombila M, Kremsner PG. Pharmacokinetics of two paediatric artesunate mefloquine drug formulations in the treatment of uncomplicated falciparum malaria in **Gabon**. *J Antimicrob Chemother.* 2007 Nov;60(5):1091-6.

Agomo PU, Meremikwu MM, Wabila IM, Omalu II, Odey FA, Oguche S, Ezeiru VI, Aina OO. Efficacy, safety and tolerability of artesunate-mefloquine in the treatment of uncomplicated Plasmodium falciparum malaria in four geographic zones of **Nigeria**. *Malar J.* 2008 Sep 9;7:172.

- Sowunmi A, Oduola AMJ, Ogundahunsi OAT, Fehintola FA, Ilesanmi OA, Akinyinka OO, Arowojolu AO. Randomised trial of artemether versus artemether and mefloquine for the treatment of chloroquine/sulfadoxine-pyrimethamine-resistant falciparum malaria during pregnancy. **Nigeria**. *Journal of Obstetrics & Gynaecology*; Jul 1998, Vol. 18 Issue 4, p322-327
- Pillai DR, Hajar G, Montoya Y, Marouñño W, Ruebush TK 2nd, Wongsrichanalai C, Kain KC. Lack of prediction of mefloquine and mefloquine-artesunate treatment outcome by mutations in the *Plasmodium falciparum* multidrug resistance 1 (*pfmdr1*) gene for *P. falciparum* malaria in **Peru**. *Am J Trop Med Hyg*. 2003 Jan;68(1):107-10.
- Grande T, Bernasconi A, Erhart A, Gamboa D, Casapia M, Delgado C, Torres K, Fanello C, Llanos-Cuentas A, D'Alessandro U. A randomised controlled trial to assess the efficacy of dihydroartemisinin-piperaquine for the treatment of uncomplicated falciparum malaria in **Peru**. *PLoS One*. 2007 Oct 31;2(10):e1101.
- Marquino W, Huilca M, Calampa C, Falconí E, Cabezas C, Naupay R, Ruebush TK 2nd. Efficacy of mefloquine and a mefloquine-artesunate combination therapy for the treatment of uncomplicated *Plasmodium falciparum* malaria in the Amazon Basin of **Peru**. *Am J Trop Med Hyg*. 2003 May;68(5):608-12.
- Gomez L EA, Jurado MH, Cambon N. Randomised efficacy and safety study of two 3-day artesunate rectal capsule/mefloquine regimens versus artesunate alone for uncomplicated malaria in **Ecuadorian** children. *Acta Tropica* 89 (2003) 47–53
- Avila JC, Villaroel R, Marquino W, Zegarra J, Mollinedo R, Ruebush TK. Efficacy of mefloquine and mefloquine-artesunate for the treatment of uncomplicated *Plasmodium falciparum* malaria in the Amazon region of **Bolivia**. *Trop Med Int Health*. 2004 Feb;9(2):217-21.
- Gutman J , Green M , Durand S , Rojas OV , Ganguly B , Quezada WM , Utz GC , Slutsker L , Ruebush TK II , Bacon DJ. Mefloquine pharmacokinetics and mefloquine-artesunate effectiveness in **Peruvian** patients with uncomplicated *Plasmodium falciparum* malaria . *Malar J* 2009 8: 58 .
- Alvarez G, Tobón A, Piñeiros JG, Ríos A and Blair S. Dynamics of *Plasmodium falciparum* Parasitemia Regarding Combined Treatment- Regimens for Acute Uncomplicated Malaria, Antioquia, **Colombia**. *Am. J. Trop. Med. Hyg.*, 83(1), 2010, pp. 90–96
- Macedo de Oliveira A, Chavez J , Ponce de Leon G , Durand S , Arrospide N, Roberts J, Cabezas C and Marquino W. Efficacy and Effectiveness of Mefloquine and Artesunate Combination Therapy for Uncomplicated *Plasmodium falciparum* Malaria in the **Peruvian** Amazon. *Am. J. Trop. Med. Hyg.*, 85(3), 2011, pp. 573–578